

# Low-mass spectroscopic binaries in the Hyades: a candidate brown dwarf companion

I. Neill Reid<sup>1</sup> and S. Mahoney<sup>2</sup>

<sup>1</sup>*Dept. of Physics & Astronomy, University of Pennsylvania, 209 S. 33rd Street, Philadelphia, PA 19104-6396;*

*e-mail: inr@morales.physics.upenn.edu*

<sup>2</sup>*Hinman Box 1709, Dartmouth College, Hanover, NH 03755*

1 February 2008

## ABSTRACT

We have used the HIRES echelle spectrograph in the Keck I telescope to obtain high-resolution spectroscopy of 51 late-type M dwarfs in the Hyades cluster. Cross-correlating the calibrated data against spectra of white dwarfs allows us to determine heliocentric velocities with an accuracy of  $\pm 0.3 \text{ kms}^{-1}$ . Twenty seven stars were observed at two epochs in 1997; two stars, RHy 42 and RHy 403, are confirmed spectroscopic binaries. RHy 42 is a double-lined, equal-mass system; RHy 403 is a single-lined, short-period binary,  $P \sim 1.275$  days. RHy 403A has an absolute magnitude of  $M_I = 10.85$ , consistent with a mass of  $0.15 M_\odot$ . The systemic mass function has a value  $\frac{(M_2 \sin(i))^3}{(M_1 + M_2)^2} = 0.0085$ , which, combined with the non-detection of a secondary peak in the cross-correlation function, implies  $0.095 > M_2 > 0.06 M_\odot$ , and the strong possibility that the companion is the first Hyades brown dwarf to be identified. Unfortunately, the maximum expected angular separation in the system is only  $\sim 0.25 \text{ mas}$ .

Five other low-mass Hyads are identified as possible spectroscopic binaries, based either on repeat observations or comparison between the observed radial velocity and the value expected for Hyades cluster members. Combined with HST imaging data, we infer a binary fraction between 23 and 30%.

All of the stars are chromospherically active. RHy 281 was caught in mid-flare and, based on that detection, we estimate a flaring frequency of  $\sim 2.5\%$  for low-mass Hyades stars. Nine stars have rotational velocities,  $v \sin(i)$ , exceeding  $20 \text{ kms}^{-1}$  and most of the sample have detectable rotation. We examine the  $H\alpha$  emission characteristics of low-mass cluster members and show that there is no evidence for a correlation with rotation.

**Key words:** stars: chromospheres, late-type, low-mass, brown dwarfs

## 1 INTRODUCTION

The frequency of binary and multiple systems amongst stars of different masses is a fundamental parameter in both star formation theory, and understanding mechanisms for the formation of planetary systems. Low-mass stars have received less attention than solar-type G dwarfs, but the available observations suggest that the multiplicity fraction decreases from  $\sim 70\%$  or more at  $M \sim 1 M_\odot$  (Duquennoy & Mayor, 1991) to  $\sim 35\%$  for M dwarfs (Fisher & Marcy, 1991; Reid & Gizis, 1997a). Most previous surveys have concentrated on field stars, which span a range of age and abundance. Open clusters offer an alternative laboratory, where age and abundance are both homogeneous and, within the

limits of theoretical models, known factors. Cluster analyses, however, must allow for the possibility that dynamical evolution may have modified key characteristics since formation. Binaries, for example, are likely to be retained preferentially, inflating the apparent binary frequency at a given spectral type.

The Hyades is the nearest substantial open cluster, and its members have received considerable observational and theoretical attention. Higher-mass cluster members (spectral types earlier than mid-K) have been the subject of both radial velocity observations (Griffin et al, 1988) and high-resolution infrared speckle interferometry (Patience et al, 1998), as well as extensive proper-motion surveys capable of detecting wide companions (Hanson, 1975; Reid, 1992 and

refs within). The deduced binary frequency is broadly consistent with Duquennoy & Mayor's analysis of similar-mass stars in the Solar Neighbourhood. At lower masses, over fifty of the cluster M dwarfs were targeted for observation by the Hubble Space Telescope (Gizis & Reid, 1997; Reid & Gizis, 1997b), with the Planetary Camera images providing a spatial resolution comparable to the ground-base speckle data,  $\Delta \sim 0.1$  arcsec. Again, the frequency of companions is consistent with that amongst nearby field dwarfs, although only six binaries are available for this statistical comparison.

Few accurate radial velocity measurements exist for late-type Hyades dwarfs. Griffin et al include a few late-K, early-M dwarfs in their survey, and Cochran & Hatzes (1999) include early-type M dwarfs in their planet search. Stauffer et al (1997) surveyed 49 early-type M dwarfs, including three non-members, identifying seven double-lined spectroscopic binaries and one triple system. The faintest star in their sample, vA 127 (RHy 143) has spectral type M4, and an absolute magnitude  $M_V = 12.83$ . Data for later type stars are scarce. Jones et al (1996) include three of the lowest luminosity cluster members (Br 804=RHy 386, Br 816 and Br 262=LP415-20, from Bryja et al, 1993) in their high-resolution spectroscopy of low-mass Hyads and Pleiads. Most recently, Terndrup et al (2000) have observed 18 Hyades dwarfs with  $M_V > 11$ , including ten in common with the current sample\*.

Given the paucity of observations, it is not surprising that few spectroscopic binaries have been identified amongst the cluster M dwarfs. Extending observations to cover this smaller range of physical separations provides a further comparison of the binary frequency relative to that amongst field dwarfs. In addition, high resolution spectroscopic data provide a means of probing the physical structure of these stars, notably the range of chromospheric activity and the distribution of rotational velocities. The current scarcity of observations reflects the faint apparent magnitudes of these stars, and the consequent difficulties of obtaining high signal-to-noise data at the appropriate spectral resolution. Those circumstances have changed with the availability of 8-10 metre class telescopes equipped with high efficiency echelle spectrographs.

This paper presents the first results from a survey of late-type Hyades M dwarfs, based on observations with the HIRES spectrograph on the Keck I telescope. Most of the targets also have high spatial-resolution HST images. The following section outlines the observations; section 3 describes the measurement of radial and rotational velocities; section 4 presents the techniques used to identify possible binaries; section 5 discusses the binaries identified from our observations; section 6 compares rotation and activity in these low mass stars; and section 7 presents our conclusions.

## 2 OBSERVATIONS

We have obtained high resolution spectroscopy of 53 late-type members of the Hyades cluster using the HIRES echelle spectrograph (Vogt et al, 1995) on the Keck I telescope.

\* Note that the stars designated Re by Terndrup et al should be identified using the prefix RHy.

The observations were made in the course of three allocations: a one night run, January 13 (UT), 1997; over the three night period December 10 to 12 in the same year; and on December 29 and 30, 1999. Conditions were clear and photometric throughout, with sub-arcsecond seeing. All of the target stars except G8-17 are drawn from Reid's (1993) list of stars with both proper motions and photometry consistent with Hyades membership. G 8-17 is listed as a candidate Hyades member in the Lowell survey (Giclas et al, 1972), and the star has UBV photometry by Sandage & Kowal (1986). Our spectrum confirms it as a Hyades member. The majority of these stars were included in Reid & Gizis' (1997b) HST Planetary Camera imaging programme, which embraced Hyades members with  $M_V > 11.9$  (spectral type later than M3). Several are therefore known to be spatially-resolved binary systems. Table 1 lists photometric data, from Reid (1993) and Leggett et al (1994), for stars in the present sample. Individual distances are estimated from the proper motions using the convergent point method and the results from Perryman et al's (1998) Hipparcos analysis (see section 4). These values are typically 5% smaller than distances cited in Reid (1993).

In addition to the Hyades members, we obtained observations of field M dwarfs of comparable spectral types, as well as several candidate Hyads which prove to be non-members. Photometry and spectral types for those stars are listed in Table 2, together with heliocentric velocity measurements, both from the published literature (Marcy & Benitz, 1989; Delfosse et al, 1998) and from our own observations. One of the Hyades non-members, RHy 80, is a double-lined spectroscopic binary, Table 2 lists velocity measurements at two epochs.

Our observations were made using a  $0''.86$  second slit, which gives a resolution of 45,000, or  $6.67 \text{ km s}^{-1}$ . With an echelle angle of  $-0^\circ 18'$  and a cross-disperser angle of  $1^\circ 37'$ , the spectra provide partial coverage of the wavelength region from 6350 Å and 8730 Å in 16 orders, including H $\alpha$  and the atmospheric A and B bands. We adopted a standard exposure time of 600 seconds for stars with  $13 < V < 17$ , increasing to integration time to between 900 and 1800 seconds for fainter stars.

The images were flat-field calibrated and bias subtracted in the usual manner, and the spectral orders extracted using software written by T. Barlow. The wavelength calibration was determined from thorium-argon lamp exposures made at the start of each night, using standard IRAF routines to apply that calibration to the individual observations. Previous experience has shown that this technique provides velocities accurate to  $\pm 1 \text{ km s}^{-1}$ . Interspersing calibration arcs between programme observations does not necessarily lead to improve the stability of the wavelength scale. However, other means are available to establish the velocity zeropoint with higher accuracy, as described in the following section. We have not attempted to set any of the spectra on an absolute flux scale.

## 3 RADIAL VELOCITY AND STELLAR ROTATION MEASUREMENTS

The primary aim of this programme is to identify spectroscopic binaries in the Hyades cluster. In general, achieving

**Table 1.** Hyades cluster members

| RHy   | Other       | V     | (V-I) | $M_V$ | r (pc.) | Comments                                       |
|-------|-------------|-------|-------|-------|---------|------------------------------------------------|
| 9     | LP 357-86   | 16.73 | 3.18  | 13.45 | 45.3    | unresolved HST                                 |
| 23    | LP 414-30   | 16.18 | 3.01  | 12.74 | 48.8    | unresolved                                     |
| 42    | LP 414-794  | 15.30 | 2.93  | 12.04 | 44.9    | unresolved                                     |
| 46    | LP 414-51   | 16.68 | 3.12  | 13.09 | 52.4    | unresolved                                     |
| 49AB  | LP 414-50   | 15.89 | 3.04  | 12.51 | 47.5    | HST binary, $\Delta = 0''.36$ , $\delta I$ 1.6 |
| 60    | LP 414-110  | 16.12 | 3.14  | 12.50 | 52.9    | unresolved                                     |
| 64    |             | 16.33 | 3.23  | 13.65 | 36.3    | unresolved                                     |
| 83    |             | 17.46 | 3.52  | 14.69 | 35.8    | unresolved                                     |
| 88A   |             | 13.06 | 1.90  | 9.13  | 61.2    | binary, $\Delta = 5''$                         |
| 88B   |             | 14.11 | 2.60  | 10.18 | 61.2    |                                                |
| 98    | LP 414-1570 | 16.63 | 3.22  | 13.64 | 36.8    | unresolved                                     |
| 101   | LP 474-1606 | 17.11 | 3.46  | 14.20 | 37.9    | unresolved                                     |
| 115   |             | 16.50 | 3.03  | 12.74 | 56.4    | unresolved                                     |
| 119AB | LP 414-138  | 17.29 | 3.54  | 14.16 | 42.3    | HST binary, $\Delta = 0''.88$ , $\delta I$ 0.1 |
| 126   | LP 414-1895 | 17.33 | 3.38  | 13.55 | 37.8    | unresolved                                     |
| 129   | LP 474-217  | 15.33 | 2.86  | 12.40 | 38.5    | unresolved                                     |
| 132   | vA 94       | 15.83 | 3.03  | 12.30 | 50.8    | unresolved                                     |
| 143   | vA 127      | 16.23 | 3.02  | 12.83 | 37.9    | unresolved                                     |
| 158   | LP475-14    | 16.31 | 3.20  | 13.07 | 44.5    | unresolved                                     |
| 162   |             | 16.54 | 3.21  | 13.63 | 38.2    | unresolved                                     |
| 163   | vA 213      | 15.44 | 3.00  | 12.48 | 39.1    | unresolved                                     |
| 165   | LP 415-19   | 19.67 | 4.34  | 16.41 | 44.9    | unresolved                                     |
| 182   | LP 4156-35  | 15.89 | 3.06  | 12.53 | 47.0    | unresolved                                     |
| 199   | LP 415-50   | 16.70 | 3.22  | 13.42 | 45.3    | unresolved                                     |
| 200   | LP 475-59   | 16.07 | 2.89  | 12.56 | 50.3    | unresolved                                     |
| 202AB | LP 415-51   | 16.70 | 3.11  | 13.34 | 37.0    | HST binary, $\Delta = 0''.28$ , $\delta I$ 1.8 |
| 219   | vA 352      | 16.37 | 2.87  | 13.03 | 46.6    | unresolved                                     |
| 221AB | LP 415-71   | 15.88 | 3.10  | 12.54 | 46.7    | HST binary, $\Delta = 0''.31$ , $\delta I$ 0   |
| 230   | LP 415-881  | 18.25 | 3.42  | 14.72 | 38.7    | unresolved                                     |
| 231   | LP 415-875  | 15.89 | 3.11  | 12.49 | 38.6    | unresolved                                     |
| 240 A |             | 18.77 | 3.72  | 15.39 | 47.5    | binary, $\Delta = 2''.5$ , $\delta I$ 0        |
| 240 B |             | 18.96 | 3.82  | 15.59 | 47.5    |                                                |
| 242   | vA 432      | 15.91 | 2.92  | 12.55 | 47.0    | unresolved                                     |
| 244AB | LP 415-108  | 15.64 | 2.97  | 12.07 | 51.8    | HST binary, $\Delta = 0''.14$ , $\delta I$ 1.1 |
| 260   | Ha 430      | 16.40 | 3.07  | 13.52 | 37.6    | unresolved                                     |
| 281   | LP 415-142  | 18.12 | 3.30  | 14.39 | 55.8    | unresolved                                     |
| 297   | LP 475-1012 | 17.04 | 3.18  | 13.42 | 52.9    | unresolved                                     |
| 298   | vA 616      | 15.90 | 2.93  | 12.66 | 44.5    | unresolved                                     |
| 301   | LP 475-125  | 16.53 | 3.15  | 13.43 | 41.6    | unresolved                                     |
| 331   | LP 415-206  | 16.43 | 2.99  | 13.30 | 42.3    | unresolved                                     |
| 346AB | LP 475-176  | 16.03 | 3.18  | 12.84 | 43.4    | HST binary, $\Delta = 0''.48$ , $\delta I$ 0.4 |
| 367   |             | 16.45 | 3.13  | 12.94 | 50.3    | unresolved                                     |
| 369   | LP 475-1419 | 16.14 | 2.92  | 12.59 | 51.3    | unresolved                                     |
| 371AB | LP 415-266  | 15.36 | 3.10  | 12.02 | 46.6    | HST binary, $\Delta = 0''.17$ , $\delta I$ 1.3 |
| 376   | LP 415-1692 | 15.80 | 2.90  | 12.52 | 45.3    | unresolved                                     |
| 377AB | LP 475-214  | 16.24 | 3.15  | 12.65 | 52.4    | binary HST, $\Delta = 1''.66$ , $\delta I$ 1.0 |
| 386   | LP 415-2128 | 18.83 | 3.69  | 15.66 | 43.0    | unresolved                                     |
| 390   | LP 415-1773 | 18.49 | 3.83  | 15.13 | 47.0    | unresolved                                     |
| 391   |             | 16.42 | 3.35  | 13.08 | 46.6    | unresolved                                     |
| 399   | LP 415-1816 | 15.09 | 2.92  | 12.18 | 38.2    | unresolved                                     |
| 402   |             | 17.77 | 3.25  | 13.94 | 58.4    | unresolved                                     |
| 403   | LP 415-327  | 18.04 | 3.47  | 14.42 | 52.9    | unresolved                                     |
|       | G8-17       | 12.68 |       | 9.3   |         | unresolved                                     |

such a goal demands multiple, high-accuracy spectroscopic observations: our initial reductions provide velocities accurate to only  $\pm 1 \text{ km s}^{-1}$ , and we have only one observation of almost half of the stars listed in Table 1. As discussed further below, we have been able to refine the accuracy of our velocity determinations. Moreover, we can utilise even

single observations for binary detection, since we have the advantage of targetting stars with known space motion.

The standard method of determining radial velocities is cross-correlation of the calibrated spectra against data for a standard star of similar spectral type and known radial velocity (Tonry & Davis, 1979). Uncertainties in the derived velocities depend both on random errors, which can be es-

**Table 2.** Field M dwarfs

| Name       | $M_V$ | (V-I) | Spectral type | $V_{hel}$<br>kms $^{-1}$ | References     | $V_{hel}$<br>kms $^{-1}$ | MJD<br>50400+ |
|------------|-------|-------|---------------|--------------------------|----------------|--------------------------|---------------|
| Gl 54.1    | 14.17 | 3.15  | M4.5          |                          | L92, RHG       | 28.0 $\pm$ 0.10          | 60.20727      |
| Gl 83.1    | 14.02 | 3.12  | M4.5          | -29                      | L92, RHG, D98  | -28.6 $^{ref}$           | 60.20251      |
|            |       |       |               |                          |                | -28.6 $^{ref}$           | 11541.17607   |
|            |       |       |               |                          |                | -28.6 $^{ref}$           | 11542.18446   |
| Gl 234 A   | 13.01 | 3.02  | M4.5          | 12                       | L92, RHG       | 14.85 $^{ref}$           | 793.59378     |
|            |       |       |               |                          |                | 16.2 $\pm$ 0.12          | 11542.46873   |
| LP 771-95  | 11.9  |       | M3.5          |                          | RHG            | -1.2 $\pm$ 0.13          | 60.21650      |
| LP 771-96A | 12.5  |       | M4            |                          | RHG            | -5.5 $\pm$ 0.18          | 60.21946      |
| Gl 402     | 12.46 | 2.79  | M4            | -1.1                     | L92, RHG, MB89 | -1.1 $\pm$ 0.20          | 792.66871     |
| Gl 445     | 12.17 | 2.64  | M4            |                          | L92, RHG       | -111.5 $\pm$ 0.25        | 794.67419     |
| Gl 447     | 13.50 | 2.98  | M4            | -31.34                   | L92, RHG, MB89 | -31.2 $\pm$ 0.24         | 794.66546     |
|            |       |       |               |                          |                | -31.3 $\pm$ 0.24         | 11541.56632   |
| Gl 831AB   | 12.50 | 2.99  | M4.5          | -57.1                    | L92, RHG, MB89 | -58.75 $\pm$ 0.30        | 793.19816     |
| Gl 876     | 11.82 | 2.74  | M4            | -1.77                    | L92, RHG, MB89 | -1.95 $\pm$ 0.25         | 793.21099     |
| RHy 80A    |       | 3.17  | M4.5          |                          | R93            | 0.88 $\pm$ 0.25          | 60.37904      |
| RHy 80B    |       | 3.2   | M4.5          |                          | R93            | 29.12 $\pm$ 0.42         |               |
| RHy 80A    |       | 3.17  | M4.5          |                          | R93            | 2.35 $\pm$ 0.25          | 792.34271     |
| RHy 80B    |       | 3.2   | M4.5          |                          | R93            | 26.4 $\pm$ 0.35          |               |
| RHy 110    |       | 3.03  | M4            |                          | R93            | 60.67 $\pm$ 0.25         | 60.42207      |
| RHy 225    |       | 2.76  | M3            |                          | R93            | 40.25 $\pm$ 0.31         | 792.54615     |
| RHy 254    |       | 3.22  | M4            |                          | R93            | -4.12 $\pm$ 0.20         | 60.25284      |

Column 6 identifies the source of the photometry, spectroscopy and radial velocity measurements listed in columns 2 to 5.

L92: photometry from Leggett, 1992

R93: photometry/spectral type from Reid, 1993

RHG: spectral type from Reid et al, 1995a

MB89: Radial velocity from Marcy & Benitz, 1989

D98: Radial velocity from Delfosse et al, 1998

timated from the width of the cross-correlation peak and inter-order agreement in the derived velocities, and on the accuracy of the wavelength calibration. In past observations, we have noted systematic differences of 1 to 1.5 kms $^{-1}$  in repeated HIRES observations of individual stars. These offsets are present in both sky and stellar spectra, even if calibrated using adjacent Th-Ar arcs. The discrepancies may originate either through differences in the illumination of the slit, or internal variations in the optical elements within the HIRES instrument.

Given those concerns, we adopt a two stage approach to determining heliocentric velocities. First, the observations are cross correlated against standard stars in the usual manner; second, we use terrestrial features in the stellar spectra to set the observations on a uniform zeropoint.

### 3.1 Relative velocity determination

In cross correlation, the spectral type of the standard template must be a good match to that of the programme stars. For the current observations, the two field stars which best satisfy this constraint are the M4.5 dwarfs Gl 83.1 (January, 1997 and December, 1999) and Gl 234A (December, 1997). Neither has published high-precision velocity measurements: heliocentric velocities for both systems have been measured by Delfosse et al (1998), but the results are cited only to the nearest kms $^{-1}$ . Moreover, Gl 234A is the primary compo-

nent in a wide binary, with a  $\sim 4.17$  AU,  $P \sim 16.60$  years,  $\frac{M_2}{M_1} \sim 0.5$  (Henry & McCarthy, 1993).

We have therefore determined heliocentric velocities for both of our template stars by cross-correlation against our data for field M dwarfs observed by Marcy & Benitz (1989): Gl 402, 447 and 876. Marcy & Benitz also observed Gl 831, but at only one epoch. Since that star is known to be a binary, it is not included in our zeropoint determination. Table 2 lists the results for all of the field stars. We derive a heliocentric velocity of -28.6 kms $^{-1}$  for Gl 83.1, with no significant variation evident in our three observations. In contrast, Gl 234A has a heliocentric velocity of 14.85 kms $^{-1}$  in December 1997, while we measure 16.17 kms $^{-1}$  in December, 1999.

The individual spectra were cross-correlated, order by order, against the appropriate template using the IRAF routine *fxcor*. The velocity differences computed by the latter routine includes heliocentric corrections. Three echelle orders are dominated by terrestrial features: 6800-6890Å, the B band; 7520-7630Å, the A-band; and 8220-8330Å. We excluded those orders when determining the mean velocity measured for the programme stars. In most cases, the remaining orders correlate well, with cross-correlation peaks of amplitude 0.7-0.9 and formal uncertainties of 0.3 to 0.7 kms $^{-1}$  in the individual velocity measurements. We have averaged the latter measurements to derive the radial velocities,  $\langle V \rangle$  listed in Tables 3 and 4, citing the rms dispersion about the mean,  $\sigma_V$ , as an estimate of the uncertainty. In

most cases, the latter values are 0.15 to 0.3  $\text{km s}^{-1}$ . Approximately 20% of the sample have substantial rotational velocities,  $v \sin(i) > 15 \text{ km s}^{-1}$ . These stars have higher uncertainties in the derived velocities.

### 3.2 Establishing the velocity zeropoint

The optimum method of wavelength calibration is to obtain a calibration spectrum simultaneously with the programme star data. Current surveys for extrasolar planets, which require accuracies exceeding  $10 \text{ m s}^{-1}$ , achieve this goal by inserting an iodide cell below the slit, superimposing iodine absorption lines on the stellar spectrum (Marcy & Butler, 1992). Unfortunately, the Hyades M dwarfs are too faint to permit use of this technique. However, Earth's atmosphere provides a ready made reference frame, superimposing a grid of  $\text{O}_2$  and  $\text{OH}$  lines on each stellar spectrum. Observations have shown that the wavelengths of these lines are stable to  $\sim 40 \text{ m s}^{-1}$ , insufficient for extrasolar-planet searches, but more than adequate for current purposes.

The terrestrial atmosphere supplies both emission and absorption lines. Of these two options, the latter is preferable for calibration purposes, since, as with the iodide cell, the optical path followed by the calibration spectrum is, perforce, identical to the stellar spectrum. In principle, we could estimate the drift in zeropoint using the Hyades/GI 83.1/GI 234A cross-correlation results for orders dominated by terrestrial absorption. However, stellar features weaken the correlation. We therefore use white dwarf stars as our terrestrial templates.

G74-7, our January 1997 template, is a DAZ white dwarf, with strong Ca II H & K absorption; HS 0507A, used for our December 1997 observations, is the brighter of a pair of DA white dwarfs detected in the course of the Hamburg Schmidt Quasar survey (Hagen et al, 1995); and 40 Eridani B, December 1999, is the brightest accessible DA white dwarf. With the exception of broad  $\text{H}\alpha$  lines, the intrinsic spectrum is a featureless continuum at red wavelengths; all absorption features are due to the Earth's atmosphere. These stars provide excellent templates for calibrating zeropoint drift of the wavelength scale.

Cross-correlating the programme stars against white dwarf spectra produces strong correlations in the orders which include the  $\text{O}_2$  A- and B-bands. Figure 1 compares the derived offsets from our January observations against the drift in velocity measured using the emission-line sky spectrum from the same observations. The agreement between the two independent measurements is good, with


$$V_{B\text{-band}} - V_{em} = 0.21 \pm 0.34 \text{ km s}^{-1}$$

and

$$V_{A\text{-band}} - V_{em} = 0.10 \pm 0.19 \text{ km s}^{-1}$$

The amplitude of variation,  $\pm 1 \text{ km s}^{-1}$ , is typical for a single night. Given these results, we have combined the A-band and B-band measurements, giving the former twice the weight of the latter, to derive the appropriate correction. Those values are listed in Table 3 as  $\Delta V_{WD}$ , and the corrected, heliocentric velocities are listed as  $V_{hel}$ .

Table 4 compares our heliocentric velocity determinations against previously published results. Setting aside RHy 158, for reasons discussed in the following section, the mean



**Figure 1.** Comparison between velocity offsets for individual spectra derived by cross-correlation using atmospheric emission lines (abscissa) and the A- and B-band absorption features. Data are plotted from January, 1997, using G74-7 as the white dwarf template.

residual, in the sense of the current measurements *minus* literature data, is

$$\Delta V_{R-O} = 0.04 \pm 0.88 \text{ km s}^{-1}, 12 \text{ stars}$$

### 3.3 Rotation

Tonry & Davis (1979) originally demonstrated that the width of the peak in the cross-correlation function is dependent on the line profiles of both template and programme object. In extragalactic astronomy, this property is used to measure velocity dispersion; in stellar astronomy, rotation is usually the dominant contributor to line profile broadening. We have used this technique to estimate rotational velocities for the stars in our sample.

The relation between the measured width of the cross-correlation peak, fitted either with a Gaussian or parabolic profile, and  $v \sin(i)$ , the projected stellar rotation, is determined using spectra of a slowly rotating star which have been broadened artificially to match known rotational velocities. Cross-correlating those spectra against a template, also with known low rotation and of similar spectral type, provides the necessary calibration. We have used our observations of GI 447 (rotation star) and GI 402 (template) to calibrate our data. Both stars have negligible rotation,  $v \sin(1) < 2.0 \text{ km s}^{-1}$  and  $< 2.3 \text{ km s}^{-1}$  respectively (Delfosse et al, 1998), and both are well matched in spectral type to the Hyades programme stars.

We chose two orders,  $\lambda\lambda 7360 - 7460 \text{ \AA}$  and  $\lambda\lambda 7840 -$

**Table 3.** Radial velocity measurements

| Star                       | $\langle V \rangle$<br>kms <sup>-1</sup> | $\sigma_V$<br>kms <sup>-1</sup> | $\Delta V_{WD}$<br>kms <sup>-1</sup> | $V_{hel}$<br>kms <sup>-1</sup> | Star        | $\langle V \rangle$<br>kms <sup>-1</sup> | $\sigma_V$<br>kms <sup>-1</sup> | $\Delta V_{WD}$<br>kms <sup>-1</sup> | $V_{hel}$<br>kms <sup>-1</sup> |
|----------------------------|------------------------------------------|---------------------------------|--------------------------------------|--------------------------------|-------------|------------------------------------------|---------------------------------|--------------------------------------|--------------------------------|
| January, 1997 field stars  |                                          |                                 |                                      |                                |             |                                          |                                 |                                      |                                |
| Gl 83.1                    | ref                                      |                                 | 0.57                                 | -28.6                          | Gl 54.1     | 28.17                                    | 0.18                            | 0.76                                 | 27.98                          |
| LP 771-95                  | -1.28                                    | 0.24                            | 0.52                                 | -1.23                          | LP 771-96AB | -5.67                                    | 0.30                            | 0.38                                 | -5.48                          |
| RHy 80                     | 1.67                                     | 0.45                            | -0.11                                | 2.35                           | RHy 110     | 59.26                                    | 0.24                            | -0.84                                | 60.67                          |
| RHy 254                    | -4.49                                    | 0.40                            | 0.20                                 | -4.12                          |             |                                          |                                 |                                      |                                |
| January, 1997 Hyads        |                                          |                                 |                                      |                                |             |                                          |                                 |                                      |                                |
| RHy 9                      | 36.82                                    | 0.28                            | 0.41                                 | 36.98                          | RHy 23      | 36.19                                    | 0.25                            | 0.33                                 | 36.43                          |
| RHy 42A                    | 30.28                                    | 0.74                            | 0.09                                 | 30.76                          | RHy 46      | 36.45                                    | 0.33                            | 0.05                                 | 36.97                          |
| RHy 42B                    | 41.70                                    | 0.81                            | 0.09                                 | 42.18                          |             |                                          |                                 |                                      |                                |
| RHy 49                     | 36.41                                    | 3.40                            | -0.25                                | 37.22                          | RHy 60      | 36.33                                    | 0.78                            | -0.10                                | 37.00                          |
| RHy 64                     | 35.71                                    | 0.86                            | -0.23                                | 36.51                          | RHy 88A     | 38.50                                    | 0.27                            | 0.00                                 | 39.07                          |
| RHy 88B                    | 33.82                                    | 0.19                            | -0.15                                | 35.04                          | RHy 98      | 38.13                                    | 3.56                            | -0.48                                | 39.19                          |
| RHy 101                    | 37.37                                    | 0.98                            | -0.28                                | 38.22                          | RHy 115     | 38.21                                    | 0.28                            | -0.13                                | 38.91                          |
| RHy 119                    | 36.70                                    | 0.41                            | -0.32                                | 37.59                          | RHy 126     | 36.18                                    | 2.61                            | -0.54                                | 37.29                          |
| RHy 129                    | 38.12                                    | 0.20                            | -0.37                                | 39.06                          | RHy 143     | 37.62                                    | 0.28                            | -0.43                                | 38.62                          |
| RHy 158                    | 38.77                                    | 0.38                            | -0.51                                | 39.85                          | RHy 163     | 36.70                                    | 7.41                            | -0.82                                | 38.09                          |
| RHy 244                    | 38.13                                    | 0.23                            | 0.74                                 | 37.96                          | RHy 260     | 40.32                                    | 0.24                            | 0.30                                 | 40.59                          |
| RHy 297                    | 39.83                                    | 0.74                            | 0.11                                 | 40.29                          | RHy 298     | 40.25                                    | 0.26                            | 0.19                                 | 40.43                          |
| RHy 301                    | 39.73                                    | 0.95                            | -0.92                                | 40.98                          | RHy 309     | 39.33                                    | 0.24                            | -0.92                                | 40.82                          |
| December, 1997 field stars |                                          |                                 |                                      |                                |             |                                          |                                 |                                      |                                |
| Gl 402                     | -4.68                                    | 0.25                            | -2.81                                | -1.10                          | Gl 445      | -113.28                                  | 0.26                            | -1.01                                | -111.50                        |
| Gl 447                     | -33.55                                   | 0.24                            | -1.60                                | -31.18                         | Gl 876      | -3.74                                    | 0.20                            | -1.13                                | -1.97                          |
| RHy 110                    | 56.11                                    | 0.09                            | -3.24                                | 60.12                          | Gl 234A     | ref                                      |                                 | 0.77                                 | 14.85                          |
| December, 1997 Hyads       |                                          |                                 |                                      |                                |             |                                          |                                 |                                      |                                |
| RHy 9                      | 33.03                                    | 0.25                            | -3.37                                | 37.17                          | RHy 23      | 32.35                                    | 0.29                            | -3.33                                | 36.45                          |
| RHy 42A                    | 14.96                                    | 1.01                            | -3.15                                | 18.88                          | RHy 46      | 32.68                                    | 0.36                            | -3.46                                | 36.91                          |
| RHy 42B                    | 53.67                                    | 0.53                            | -3.15                                | 57.59                          |             |                                          |                                 |                                      |                                |
| RHy 60                     | 32.31                                    | 0.78                            | -3.61                                | 36.69                          | RHy 64      | 31.83                                    | 0.87                            | -3.49                                | 36.09                          |
| RHy 83                     | 33.27                                    | 0.45                            | -3.62                                | 37.66                          | RHy 88A     | 35.05                                    | 0.35                            | -3.30                                | 39.13                          |
| RHy 88B                    | 32.18                                    | 0.24                            | -3.18                                | 36.13                          | RHy 98      | 32.73                                    | 1.65                            | -3.29                                | 36.89                          |
| RHy 101                    | 33.32                                    | 0.41                            | -3.65                                | 37.74                          | RHy 115     | 34.66                                    | 0.39                            | -3.63                                | 39.06                          |
| RHy 119                    | 33.91                                    | 0.37                            | -3.01                                | 37.69                          | RHy 129     | 34.96                                    | 0.39                            | -3.19                                | 38.92                          |
| RHy 143                    | 34.43                                    | 0.40                            | -3.31                                | 38.51                          | RHy 158     | 33.55                                    | 0.37                            | -3.31                                | 37.63                          |
| RHy 162                    | 34.28                                    | 2.48                            | -2.84                                | 37.89                          | RHy 163     | 34.82                                    | 0.70                            | -2.87                                | 38.46                          |
| RHy 182                    | 35.52                                    | 0.39                            | -2.98                                | 39.27                          | RHy 199     | 35.20                                    | 0.71                            | -2.84                                | 38.81                          |
| RHy 200                    | 36.27                                    | 0.52                            | -2.79                                | 39.83                          | RHy 202     | 35.19                                    | 0.38                            | -2.75                                | 38.71                          |
| RHy 219                    | 35.87                                    | 0.54                            | -2.63                                | 39.27                          | RHy 221     | 37.08                                    | 0.33                            | -2.70                                | 40.55                          |
| RHy 225                    | 36.91                                    | 0.31                            | -2.57                                | 40.25                          | RHy 231     | 35.68                                    | 0.42                            | -2.65                                | 39.10                          |
| RHy 230                    | 37.40                                    | 1.38                            | -1.51                                | 39.68                          | RHy 240A    | 37.59                                    | 1.34                            | -1.37                                | 39.73                          |
| RHy 240B                   | 36.22                                    | 0.21                            | -1.24                                | 38.33                          | RHy 260     | 38.73                                    | 0.25                            | -1.08                                | 40.68                          |
| RHy 281                    | 31.26                                    | 1.32                            | -1.29                                | 33.32                          | RHy 297     | 38.42                                    | 0.54                            | -1.14                                | 40.33                          |
| RHy 301                    | 38.27                                    | 0.54                            | -1.04                                | 40.08                          | RHy 331     | 38.23                                    | 0.30                            | -1.13                                | 40.13                          |
| RHy 346                    | 39.00                                    | 0.20                            | -1.10                                | 40.90                          | RHy 367     | 39.18                                    | 0.20                            | -1.15                                | 41.10                          |
| RHy 371                    | 36.37                                    | 0.67                            | -1.09                                | 38.23                          | RHy 377     | 40.65                                    | 0.47                            | -0.98                                | 42.40                          |
| G 8-17                     | 34.96                                    | 0.22                            | -1.12                                | 36.85                          | RHy 386     | 39.41                                    | 0.51                            | -1.10                                | 41.30                          |
| RHy 391                    | 37.88                                    | 0.82                            | -1.04                                | 39.69                          | RHy 399     | 38.97                                    | 0.25                            | -1.02                                | 40.76                          |
| RHy 402                    | 39.23                                    | 0.52                            | -0.92                                | 40.92                          | RHy 403     | 41.22                                    | 0.47                            | -0.97                                | 42.96                          |
| RHy 165                    | 36.82                                    | 2.10                            | -0.92                                | 38.51                          | RHy 390     | 39.12                                    | 1.63                            | -1.05                                | 40.94                          |
| RHy 242                    | 39.22                                    | 0.41                            | -0.46                                | 40.45                          | RHy 132     | 37.56                                    | 0.41                            | -0.50                                | 38.83                          |
| RHy 376                    | 39.90                                    | 0.32                            | -0.33                                | 41.00                          | RHy 83      | 36.79                                    | 0.52                            | -0.28                                | 37.84                          |
| RHy 46                     | 35.58                                    | 0.26                            | -0.54                                | 36.89                          | RHy 46      | 35.61                                    | 0.29                            | -0.35                                | 36.73                          |
| RHy 46                     | 35.73                                    | 0.32                            | -0.41                                | 36.91                          | RHy 230     | 38.46                                    | 1.59                            | -0.57                                | 39.80                          |
| RHy 281                    | 30.97                                    | 1.07                            | -1.06                                | 32.80                          | RHy 297     | 38.42                                    | 0.39                            | -1.22                                | 40.41                          |
| RHy 386                    | 39.60                                    | 0.79                            | -1.21                                | 41.58                          | RHy 402     | 38.95                                    | 0.78                            | -1.25                                | 40.97                          |
| RHy 403                    | 71.41                                    | 0.74                            | -1.58                                | 73.76                          |             |                                          |                                 |                                      |                                |

7960Å, for these measurements, avoiding both TiO band-heads and the stronger terrestrial features. The Gl 447 spectrum was broadened following Gray's (1976) prescription. The HIRES data have a resolution of  $R \sim 45,000$ , corresponding to a velocity of  $v_{FWHM} = 6.67 \text{ kms}^{-1}$ . This sets

a lower detection limit of  $v \sin(i) \sim 2.5 \text{ kms}^{-1}$  for our rotation measurements. Results for the full sample of Hyades stars are tabulated and discussed in section 5. Table 4 includes comparison of our measurements against previously-published results. We derive slightly lower rotational veloc-

**Table 3.** Radial velocity measurements

| Star                       | $\langle V \rangle$<br>kms $^{-1}$ | $\sigma_V$<br>kms $^{-1}$ | $\Delta V_{WD}$<br>kms $^{-1}$ | $V_{hel}$<br>kms $^{-1}$ | Star    | $\langle V \rangle$<br>kms $^{-1}$ | $\sigma_V$<br>kms $^{-1}$ | $\Delta V_{WD}$<br>kms $^{-1}$ | $V_{hel}$<br>kms $^{-1}$ |
|----------------------------|------------------------------------|---------------------------|--------------------------------|--------------------------|---------|------------------------------------|---------------------------|--------------------------------|--------------------------|
| December, 1999 field stars |                                    |                           |                                |                          |         |                                    |                           |                                |                          |
| Gl 83.1                    | ref                                |                           | 0.71                           | -28.6                    | Gl 447  | -30.63                             | 0.32                      | 1.57                           | -31.49                   |
| Gl 83.1                    | ref                                |                           | 0.0                            | -28.6                    | Gl 234A | 16.35                              | 0.12                      | 0.18                           | 16.17                    |
| Gl 447                     | -30.90                             | 0.15                      | -31.21                         | 0.31                     |         |                                    |                           |                                |                          |
| December, 1999 Hyads       |                                    |                           |                                |                          |         |                                    |                           |                                |                          |
| RHy 42                     | 36.31                              | 0.16                      | 0.95                           | 36.07                    | RHy 281 | 32.37                              | 1.19                      | 0.42                           | 32.66                    |
| RHy 403                    | 69.01                              | 0.56                      | 0.29                           | 69.42                    | RHy 42  | 35.88                              | 0.25                      | 0.0                            | 36.59                    |
| RHy 403                    | 74.68                              | 0.42                      | -0.02                          | 75.41                    | RHy 42  | 36.18                              | 0.33                      | -0.15                          | 37.04                    |
| RHy 403                    | 78.02                              | 0.34                      | -0.10                          | 78.83                    | RHy 42  | 36.23                              | 0.28                      | 0.48                           | 33.19                    |
| RHy 281                    | 32.36                              | 0.51                      | -0.21                          | 33.19                    | RHy 403 | 79.24                              | -0.04                     | 79.99                          |                          |
| RHy 42                     | 36.10                              | 0.31                      | 0.42                           | 36.39                    | RHy 403 | 78.49                              | 0.40                      | 0.74                           | 78.46                    |
| RHy 42                     | 36.74                              | 0.27                      | 0.44                           | 37.01                    | RHy 403 | 75.24                              | 0.64                      | 0.80                           | 75.15                    |
| RHy 42                     | 37.08                              | 0.31                      | 0.72                           | 37.07                    | RHy 403 | 72.48                              | 0.44                      | 1.03                           | 72.16                    |
| RHy 342                    | 36.79                              | 0.16                      | 0.04                           | 36.75                    | RHy 403 | 20.27                              | 0.46                      | 0.04                           | 20.23                    |
| RHy 42                     | 35.66                              | 0.57                      | 0.07                           | 35.59                    | RHy 403 | 27.72                              | 0.64                      | 0.03                           | 27.69                    |
| RHy 42                     | 36.39                              | 0.11                      | -0.46                          | 36.85                    | RHy 403 | 36.80                              | 0.37                      | -0.40                          | 37.20                    |
| RHy 42                     | 36.24                              | 0.17                      | -0.35                          | 36.59                    | RHy 281 | 32.02                              | 1.19                      | -0.75                          | 32.77                    |
| RHy 403                    | 46.33                              | 0.57                      | -0.71                          | 47.04                    | RHy 42  | 35.92                              | 0.28                      | -0.43                          | 36.35                    |
| RHy 403                    | 56.50                              | 0.35                      | -0.23                          | 56.73                    | RHy 42  | 36.17                              | 0.35                      | -0.42                          | 36.59                    |
| RHy 403                    | 63.94                              | 0.33                      | -0.19                          | 64.13                    |         |                                    |                           |                                |                          |

**Table 4.** Comparison with previous observations: Hyades stars

| Star     | $V_{hel}$<br>this paper | $v \sin(i)$ | $V_{hel}$<br>other | $v \sin(i)$ | reference            |
|----------|-------------------------|-------------|--------------------|-------------|----------------------|
| RHy 143  | 38.6                    | 4           | 37.5               | 6.5         | Stauffer et al, 1997 |
| RHy 225  | 40.3                    | < 2.5       | 39.2               | < 6         |                      |
| RHy 386  | 41.5                    | 9           | 41.1               | 11          |                      |
| RHy 23   | 36.4                    | 4           | 36.3               | 6.5         | Terndrup et al, 2000 |
| RHy 60   | 36.85                   | 19          | 37                 | 19          |                      |
| RHy 115  | 38.9                    | 7           | 39.0               | 9           |                      |
| RHy 119  | 37.6                    | 9           | 37.1               | 9           |                      |
| RHy 158  | 37.6/39.5               | 11          | 40.9               | 11          |                      |
| RHy 202  | 38.7                    | 2.5         | 38.8               | < 5         |                      |
| RHy 240b | 38.2                    | 14.5        | 40.5               | 14.5        |                      |
| RHy 346  | 40.9                    | 3.5         | 41.1               | 6           |                      |
| RHy 386  | 41.5                    | 9           | 41.9               | 11.5        |                      |
| RHy 390  | 40.9                    | 18          | 40.5               | 18          |                      |

ities for the slow rotators. However, the agreement is excellent for moderately and rapidly rotation stars.

#### 4 IDENTIFYING SPECTROSCOPIC BINARIES

The conventional method of identifying spectroscopic binaries is to compare velocity measurements obtained at different epochs. Table 5 lists the relevant data from our observations, including both  $V_{hel}$  and the Modified Julian Date for each observation. Almost half of the sample have only a single observation, but even those observations can be used to identify candidate spectroscopic binaries since our targets are members of the Hyades cluster.

Perryman et al (1998) have used astrometric data from the Hipparcos satellite and CORAVEL radial velocity observations to determine the mean distance of Hyades cluster

members (46.34 $\pm$ 0.27 pc.) and the cluster space motion (45.72 kms $^{-1}$ ). As noted above, those parameters correspond to a distance scale shorter by  $\sim$ 5% than that adopted in Reid (1993). In traditional open cluster analyses, absolute proper motion measurements are used to determine the location in  $(\alpha, \delta)$  of the convergent point. Given a known space motion for the cluster, the radial velocity of an idealised cluster member,  $V_{rad}^P$ , depends only on the angular distance from the convergent point,  $\lambda_{CP}$ . Similarly, the proper motion of a cluster members depends on both  $\lambda_{CP}$  and distance, allowing distance estimates to individual cluster members (Table 1).

Circumstances are more complicated for a relatively old cluster, such as the Hyades. As Perryman et al point out, the velocity dispersion of  $\sigma \sim 0.2$  kms $^{-1}$  leads to an intrinsic fuzziness in the convergent ‘point’. Thus, when we invert the process, and predict  $V_{rad}$  based on an idealised convergent point, those predictions are uncertain by  $\pm 0.2$

$\text{kms}^{-1}$ . Indeed, since the Hyades is sufficiently old to have undergone significant mass segregation, the velocity dispersion (and corresponding uncertainty in  $V_{rad}^P$ ) may be as high as  $0.3\text{--}0.4 \text{ kms}^{-1}$  for low-mass stars. Nonetheless, taking into account these uncertainties, a significant difference between the observed velocity and the expected velocity indicates either binary motion, or non-membership of the cluster. In the case of Hyades M dwarfs, chromospheric activity provides a means of deciding between those alternatives.

We have adopted the convergent point determined by Perryman et al from their maximum likelihood analysis: they derive ( $\alpha = 96.5$ ,  $\delta = 5.8$ ) from a sample of 148 Hyades members. We also use their estimate of the Hyades space motion,  $V = 45.72 \text{ kms}^{-1}$ . Table 5 lists both the expected radial velocity,  $V^P$ , for each Hyades star, and the observed-computed residual,

$$\delta V_{hel}^P = V_{hel} - V^P$$

We note that the majority of the residuals are positive, with  $\langle \delta V_{hel}^P \rangle = 0.64 \pm 0.68$  (excluding candidate binaries), suggesting that the kinematic model we have adopted for the cluster motion may require revision. The agreement is adequate for current purposes.

In order to assess the significance of differences between the predicted and observed velocities, we need to determine the uncertainty,  $\epsilon_V$ , associated with  $\delta V_{hel}^P$ . These uncertainties stem from two sources: empirical measuring accuracy, and intrinsic dispersion in cluster motions. Our data provide several means of estimating the former:


- (i) five observations in 1997 of RHy 46 give  $\langle V_{hel} \rangle = 36.88 \pm 0.09 \text{ kms}^{-1}$ ;
- (ii) thirteen observations (December, 1999) of RHy 42 give  $\langle V_{hel} \rangle = 36.57 \pm 0.42 \text{ kms}^{-1}$ ;
- (iii) five observations (1997-1999) of RHy 281 give  $\langle V_{hel} \rangle = 32.95 \pm 0.29 \text{ kms}^{-1}$ ;
- (iv) twenty-one repeat observations of twenty stars with  $|\Delta V_{epoch}| < 1 \text{ kms}^{-1}$  give a mean difference of  $-0.115 \pm 0.34 \text{ kms}^{-1}$ .

On that basis, we estimate that an rms uncertainty of  $0.35 \text{ kms}^{-1}$  is appropriate for the heliocentric velocity measurements given in Table 3. If we assume that the intrinsic velocity dispersion of Hyades M dwarfs has a similar value, then we derive a minimum value of  $\epsilon_V = 0.5 \text{ kms}^{-1}$ . If  $\sigma_V$ , the order-to-order rms dispersion of velocity measurements for a given observation, exceeds  $0.5 \text{ kms}^{-1}$  (Table 3), then we let  $\epsilon_V = \sigma_V$ . Column 7 in Table 4 gives the ratio between  $\delta V_{hel}^P$  and  $\epsilon_V$  for each star.

## 5 DISCUSSION

### 5.1 New spectroscopic binaries

Figure 2 plots the colour-magnitude diagram for lower main-sequence in the Hyades, marking the spectroscopic binaries and candidate binaries identified from our observations. Three stars in Table 4 are prime binary candidates: RHy 42, 281 and 403. Two are confirmed as binaries based on the 1997 data alone: RHy 42, a double-lined system; and RHy 403, a single-lined system, whose radial velocity increased by  $31 \text{ kms}^{-1}$  in only 26 hours in December, 1997. The third



**Figure 2.** The  $(M_V, (V-I))$  colour-magnitude diagram for late-type Hyades dwarfs. Crosses are stars lacking HST observations; open triangles are stars unresolved by HST (RG97); solid triangles are resolved HST binaries (RG97); solid points mark the locations of RHy 42, 281 and 403; the three possible velocity variables, RHy 158, 221 and 377, are identified using open circles, while open squares identify four stars with unusual cross-correlation profiles: RHy 49, 244, 162 and 165.

star, RHy 281, exhibits no variation in velocity, but the measured value is offset by  $\sim 5\sigma$  from the expected value for a cluster member. These three stars were therefore chosen as the focus of our most recent observations,

Figure 3 plots the results of our December, 1999 observations. RHy 403 is confirmed as a spectroscopic binary. The curve plotted is a simple sinewave with a period of 1.276 days and an amplitude of  $40 \text{ kms}^{-1}$ , matched by eye to the data. Phase zero is at MJD=51541.3521 days. The uncertainty in the derived period is  $\pm 0.005$  days. The December 1997 observations are consistent with being obtained at phases of  $\phi = 0.487$  and  $\phi = 0.339$  respectively, with  $\phi = 0.25$  at 40794.429. The total number of cycles between the latter date and the observed maximum on December 29 can only be given as  $584 \pm 2$ .

The cross-correlation functions from our 1997 observations of RHy 42 clearly identify the star as a double-lined spectroscopic binary, as illustrated in figure 4. We also plot  $H\alpha$  emission line data: the unusual profile in December, 1997 is the result of the superposition of two lines of almost equal strength, each possessing a central reversal. The Ca I line at  $6598\text{\AA}$  is also clearly double at the latter epoch. However, there is no evidence for either velocity variations or line-splitting in the December, 1999 data.

These observations are consistent with the identification



**Table 5.** Heliocentric velocities

| RHy              | MJD<br>50000+ | $V_{hel}$<br>kms <sup>-1</sup> | $\epsilon_V$<br>kms <sup>-1</sup> | $V^P$<br>kms <sup>-1</sup> | $\delta V_{hel}^P$<br>kms <sup>-1</sup> | $\delta V/\epsilon_V$ | $\Delta V_{epoch}$<br>kms <sup>-1</sup> | H $\alpha$ EW<br>Å |
|------------------|---------------|--------------------------------|-----------------------------------|----------------------------|-----------------------------------------|-----------------------|-----------------------------------------|--------------------|
| 9                | 460.31252     | 36.98                          | 0.5                               | 36.12                      | 0.86                                    | 1.7                   |                                         | 5.1                |
|                  | 792.24254     | 37.22                          | 0.5                               |                            | 1.1                                     | 2.2                   | 0.24                                    | 3.8                |
| 23               | 460.32099     | 36.42                          | 0.5                               | 35.70                      | 0.72                                    | 1.4                   |                                         | 3.4                |
|                  | 792.25098     | 36.50                          | 0.5                               |                            | 0.8                                     | 1.6                   | 0.08                                    | 4.4                |
| 42               | 460.32924     | A 30.77                        | 0.7                               | 36.19                      |                                         |                       |                                         | 3.4                |
|                  |               | B 42.18                        | 0.8                               |                            | 0.28                                    | 0.4                   |                                         |                    |
|                  | 792.25906     | A 18.88                        | 0.6                               |                            |                                         |                       |                                         | 4.3                |
|                  |               | B 57.59                        | 0.5                               |                            | 2.0                                     | 4                     |                                         |                    |
|                  | 1541.18103    | 36.07                          | 0.5                               |                            | -0.12                                   | -0.2                  |                                         | 3.5                |
|                  | 1541.23980    | 36.59                          | 0.5                               |                            | 0.40                                    | 0.8                   |                                         | 4.0                |
|                  | 1541.29145    | 37.04                          | 0.5                               |                            | 0.75                                    | 1.5                   |                                         | 3.8                |
|                  | 1541.33793    | 36.46                          | 0.5                               |                            | 0.27                                    | 0.5                   |                                         | 3.8                |
|                  | 1541.40453    | 36.39                          | 0.5                               |                            | 0.20                                    | 0.4                   |                                         | 3.7                |
|                  | 1541.44922    | 37.01                          | 0.5                               |                            | 0.82                                    | 1.6                   |                                         | 3.5                |
|                  | 1541.47928    | 37.07                          | 0.5                               |                            | 0.86                                    | 1.7                   |                                         | 3.9                |
|                  | 1542.18976    | 36.75                          | 0.5                               |                            | 0.56                                    | 1.1                   |                                         | 4.7                |
|                  | 1542.24088    | 35.59                          | 0.5                               |                            | -0.60                                   | 1.2                   |                                         | 4.9                |
|                  | 1542.28896    | 36.85                          | 0.5                               |                            | 0.66                                    | 1.3                   |                                         | 5.6                |
|                  | 1542.33138    | 36.59                          | 0.5                               |                            | 0.40                                    | 0.8                   |                                         | 4.7                |
|                  | 1542.38990    | 36.35                          | 0.5                               |                            | 0.16                                    | 0.3                   |                                         | 4.6                |
|                  | 1542.43474    | 36.59                          | 0.5                               |                            | 0.40                                    | 0.8                   |                                         | 4.7                |
| 46               | 460.34582     | 36.97                          | 0.5                               | 36.19                      | 0.78                                    | 1.6                   |                                         | 3.8                |
|                  | 792.31009     | 36.91                          | 0.5                               |                            | 0.72                                    | 1.4                   | -0.06                                   | 3.9                |
|                  | 794.39999     | 36.90                          | 0.5                               |                            | 0.73                                    | 1.4                   | -0.07                                   | 4.1                |
|                  | 794.40812     | 36.73                          | 0.5                               |                            | 0.54                                    | 1.1                   | -0.24                                   | 4.2                |
|                  | 794.41615     | 36.91                          | 0.5                               |                            | 0.72                                    | 1.4                   | -0.06                                   | 4.2                |
|                  | 460.34517     | 37.22                          | 3.4                               | 36.02                      | 1.2                                     | 0.4                   |                                         | 4.8                |
| 49 <sup>r</sup>  | 792.31824     | 34.75                          | 2.0                               |                            | -1.27                                   | -0.7                  | -2.47                                   | 4.4                |
|                  | 460.36269     | 37.00                          | 0.8                               | 36.43                      | 0.57                                    | 0.7                   |                                         | 3.1                |
| 60               | 792.32645     | 36.70                          | 0.8                               |                            | 0.27                                    | 0.3                   | -0.3                                    | 3.9                |
|                  | 460.37087     | 36.23                          | 0.9                               | 36.33                      | -0.1                                    | -0.1                  |                                         | 3.2                |
| 64               | 792.33456     | 36.09                          | 0.7                               |                            | -0.24                                   | -0.3                  | -0.14                                   | 3.4                |
|                  | 792.35098     | 37.66                          | 0.5                               | 37.14                      | 0.52                                    | 1.0                   |                                         | 4.8                |
| 83               | 794.42461     | 37.84                          | 0.5                               |                            | 0.70                                    | 1.4                   | 0.18                                    | 3.9                |
|                  | 460.39414     | 39.07                          | 0.5                               | 36.61                      | 2.46                                    | 4.9                   |                                         | 0.5a               |
| 88A              | 792.35970     | 39.13                          | 0.5                               |                            | 2.52                                    | 5.0                   | 0.06                                    | 0.5a               |
|                  | 460.40029     | 35.04                          | 0.5                               | 36.61                      | -1.57                                   | -3.2                  |                                         | 2.8                |
| 88B              | 792.36787     | 36.13                          | 0.5                               |                            | -0.48                                   | 1.0                   | 1.09                                    | 1.4                |
|                  | 460.40542     | 39.19                          | 3.6                               | 36.77                      | 2.42                                    | 0.7                   |                                         | 5.0                |
| 98 <sup>r</sup>  | 792.37262     | 36.91                          | 1.65                              |                            | 0.14                                    | 0.1                   | -2.28                                   | 3.7                |
|                  | 460.41391     | 38.22                          | 1.0                               | 37.85                      | 0.37                                    | 0.4                   |                                         | 4.3                |
| 101              | 792.38320     | 37.74                          | 0.5                               |                            | -0.11                                   | -0.2                  | -0.48                                   | 4.4                |
|                  | 460.43029     | 38.91                          | 0.5                               | 37.32                      | 1.59                                    | 3.2                   |                                         | 3.6                |
| 115              | 792.40683     | 39.06                          | 0.5                               |                            | 1.74                                    | 3.5                   | 0.17                                    | 3.5                |
|                  | 460.43856     | 37.59                          | 0.5                               | 37.14                      | 0.45                                    | 0.9                   |                                         | 4.4                |
| 119              | 792.41942     | 37.68                          | 0.5                               |                            | 0.54                                    | 1.1                   | 0.09                                    | 4.0                |
|                  | 460.44699     | 37.29                          | 2.6                               | 37.76                      | -0.47                                   | -0.2                  |                                         | 6.0                |
| 126 <sup>r</sup> | 460.45539     | 39.06                          | 0.5                               | 38.04                      | 1.02                                    | 2                     |                                         | 3.6                |
|                  | 792.42803     | 38.92                          | 0.5                               |                            | 0.88                                    | 1.8                   | -0.14                                   | 4.0                |
| 129              | 793.50806     | 38.83                          | 0.5                               | 37.81                      | 1.02                                    | 2.0                   |                                         | 4.3                |

of RHy 42 as a moderate-period binary system. The semi-major axis in an equal-mass binary is given by

$$a = \frac{\mathcal{G}(M_1 + M_2)}{v_{circ}^2}$$

Both components have  $M_V \sim 12.5$ , or, from Henry & McCarthy's (1993) (mass- $M_V$ ) relation,  $M \sim 0.2M_\odot$ . If we assume that the observed velocity difference in December, 1997,  $\delta V = 34$  kms<sup>-1</sup>, is close to the maximum value, and an inclination of 45°, then  $v_{circ} \sim 24$  kms<sup>-1</sup> and  $P \sim 0.75$

years. Presumably our 1999 observations happened to catch the system at phase  $\phi = 0^\circ$  or  $180^\circ$ , with both stars having radial velocities close to the systemic value.

Although RHy 281 has a radial velocity offset by 6 kms<sup>-1</sup> from that expected for a Hyades star, our data show no evidence for significant velocity variation. The rms dispersion of all five observations, spanning two years, is only 0.29 kms<sup>-1</sup>. The average H $\alpha$  emission properties of this star are entirely consistent with Hyades membership. It is possi-

**Table 5.** Heliocentric velocities (contd.)

| RHy              | MJD<br>50000+ | $V_{hel}$<br>kms <sup>-1</sup> | $\epsilon_V$<br>kms <sup>-1</sup> | $V^P$<br>kms <sup>-1</sup> | $\delta V_{hel}^P$<br>kms <sup>-1</sup> | $\delta V/\epsilon_V$ | $\Delta V_{epoch}$<br>kms <sup>-1</sup> | H $\alpha$ EW<br>Å |
|------------------|---------------|--------------------------------|-----------------------------------|----------------------------|-----------------------------------------|-----------------------|-----------------------------------------|--------------------|
| 143              | 460.46358     | 38.62                          | 0.5                               | 37.89                      | 0.73                                    | 1.5                   |                                         | 3.0                |
|                  | 792.43758     | 38.51                          | 0.5                               |                            | 0.62                                    | 1.2                   | -0.11                                   | 6.6                |
| 158              | 460.47216     | 39.45                          | 0.5                               | 38.71                      | 0.74                                    | 1.5                   |                                         | 4.1                |
|                  | 792.44589     | 37.63                          | 0.5                               |                            | -1.08                                   | -2.2                  | -1.82                                   | 4.2                |
| 162 <sup>r</sup> | 460.48031     | 38.09                          | 5.0                               | 37.83                      | 0.26                                    | 0.05                  |                                         | 4.7                |
|                  | 792.47709     | 37.87                          | 2.5                               |                            | 0.04                                    | 0                     | -0.22                                   | 5.3                |
| 163              | 792.48866     | 38.46                          | 0.7                               | 38.30                      | 0.16                                    | 0.2                   |                                         | 3.2                |
| 165 <sup>r</sup> | 793.47003     | 38.51                          | 2.1                               | 37.82                      | 0.69                                    | 0.3                   |                                         | 7.7                |
| 165 <sup>r</sup> | 793.47003     | 38.51                          | 2.1                               | 37.82                      | 0.69                                    | 0.3                   |                                         | 7.7                |
| 182              | 792.49718     | 39.27                          | 0.5                               | 38.59                      | 0.68                                    | 1.4                   |                                         | 4.0                |
| 199              | 792.50523     | 38.81                          | 0.7                               | 38.26                      | 0.55                                    | 0.7                   |                                         | 4.1                |
| 200              | 792.51342     | 39.83                          | 0.5                               | 39.00                      | 0.83                                    | 1.6                   |                                         | 3.6                |
| 202              | 792.52180     | 38.71                          | 0.5                               | 38.31                      | 0.4                                     | 0.8                   |                                         | 3.0                |
| 219              | 792.52990     | 39.27                          | 0.5                               | 38.73                      | 0.54                                    | 1.1                   |                                         | 4.3                |
| 221              | 792.53800     | 40.55                          | 0.5                               | 38.63                      | 1.92                                    | 3.8                   |                                         | 2.9                |
| 230              | 793.24775     | 39.68                          | 1.4                               | 38.74                      | 0.94                                    | 0.7                   |                                         | 5.2                |
|                  | 794.44496     | 39.80                          | 1.6                               |                            | 1.06                                    | 0.7                   | 0.12                                    | 4.2                |
| 231              | 792.55084     | 39.10                          | 0.5                               | 38.63                      | 0.47                                    | 0.9                   |                                         | 5.8                |
| 240A             | 793.26704     | 39.73                          | 1.3                               | 38.90                      | 0.83                                    | 0.6                   |                                         | 6.0                |
| 240B             | 793.27862     | 38.23                          | 0.5                               | 38.90                      | -0.67                                   | 1.3                   |                                         | 4.0                |
| 242              | 793.50000     | 40.45                          | 0.5                               | 39.16                      | 1.26                                    | 2.5                   |                                         | 3.1                |
| 244              | 460.24334     | 37.96                          | 0.5                               | 38.54                      | -0.58                                   | -1.2                  |                                         | 4.7                |
| 260              | 460.26471     | 40.59                          | 0.5                               | 39.28                      | 1.31                                    | 2.6                   |                                         | 3.6                |
|                  | 793.28678     | 40.58                          | 0.5                               |                            | 1.30                                    | 2.6                   | -0.01                                   | 4.3                |
| 281              | 793.29503     | 33.32                          | 1.3                               | 39.22                      | -5.80                                   | -4.5                  |                                         | 4.6                |
|                  | 794.45771     | 32.80                          | 1.1                               |                            | -6.42                                   | -5.8                  | -0.52                                   | 22.8               |
|                  | 1541.18724    | 32.66                          | 1.2                               |                            | -6.56                                   | -5.5                  | -0.66                                   | 6.7                |
|                  | 1541.34444    | 33.19                          | 0.6                               |                            | -6.03                                   | -10.1                 | -0.13                                   | 6.1                |
|                  | 1542.33699    | 32.77                          | 1.2                               |                            | -6.45                                   | -5.4                  | -0.55                                   | 5.5                |
| 297              | 460.27310     | 40.29                          | 0.5                               | 39.66                      | 0.63                                    | 1.3                   |                                         | 4.1                |
|                  | 793.30331     | 40.33                          | 0.5                               |                            | 0.67                                    | 1.3                   | 0.04                                    | 4.7                |
|                  | 794.49131     | 40.41                          | 0.5                               |                            | 0.75                                    | 1.5                   | 0.12                                    | 4.3                |
| 298              | 460.28963     | 40.63                          | 0.5                               | 39.68                      | 1.05                                    | 2.1                   |                                         | 3.3                |
|                  | 793.31490     | 40.61                          | 0.5                               |                            | 1.03                                    | 2.1                   | -0.02                                   | 4.1                |
| 301              | 460.48851     | 40.98                          | 1.0                               | 39.77                      | 1.2                                     | 1.2                   |                                         | 5.0                |
|                  | 793.32307     | 40.08                          | 0.5                               |                            | 0.31                                    | 0.6                   | -0.9                                    | 6.1                |
| 309              | 460.49697     | 40.82                          | 0.5                               | 39.15                      | 1.67                                    | 3.1                   |                                         | 4.9                |
| 331              | 793.33144     | 40.13                          | 0.5                               | 39.28                      | 0.85                                    | 1.7                   |                                         | 3.0                |
| 346              | 793.34021     | 40.90                          | 0.5                               | 40.16                      | 0.74                                    | 1.5                   |                                         | 3.5                |
| 367              | 793.34828     | 41.10                          | 0.5                               | 40.24                      | 0.86                                    | 1.7                   |                                         | 4.8                |
| 369              | 793.35632     | 41.51                          | 0.5                               | 40.30                      | 1.21                                    | 2.4                   |                                         | 2.7                |
| 371              | 793.36490     | 38.23                          | 0.7                               | 39.82                      | -1.59                                   | -2.3                  |                                         | 3.8                |
| 376              | 793.51613     | 41.00                          | 0.5                               | 39.99                      | 1.01                                    | 2.0                   |                                         | 3.8                |
| 377              | 793.37344     | 42.40                          | 0.5                               | 40.38                      | 2.02                                    | 4.0                   |                                         | 3.8                |
| 386              | 793.40658     | 41.30                          | 0.5                               | 40.08                      | 1.22                                    | 2.4                   |                                         | 4.4                |
|                  | 794.50953     | 41.58                          | 0.8                               |                            | 1.50                                    | 1.9                   | 0.28                                    | 4.8                |
| 390              | 793.48502     | 40.94                          | 1.6                               | 39.84                      | 1.10                                    | 0.7                   |                                         | 8.7                |
| 391              | 793.42219     | 39.69                          | 0.8                               | 39.61                      | 0.08                                    | 0.1                   |                                         | 5.1                |
| 399              | 793.43026     | 40.76                          | 0.5                               | 39.90                      | 0.86                                    | 1.7                   |                                         | 3.8                |
| 402              | 793.43834     | 40.92                          | 0.5                               | 39.92                      | 1.0                                     | 2.0                   |                                         | 2.9                |
|                  | 794.53181     | 39.97                          | 0.8                               |                            | 0.05                                    | 0                     | -0.95                                   | 3.8                |


ble that RHy 281 may be no longer be bound to the Hyades cluster proper, although still part of the more extensive moving group. Alternatively, RHy 281 may be a long-period binary system, with an as-yet undetected low-mass companion. Further observations of this star are clearly warranted.

Seven other stars are identified as candidate binaries: three on the basis of the velocity data listed in table 4; four display anomalies in the shape of the cross-correlation peak,

with one star almost certainly binary. In the former group, RHy 221 and 377 both have only a single velocity determination, but in both cases,  $\delta V/\epsilon_V \sim 4$ . Both are known as a binary from HST observations. The peak in RHy 377's cross-correlation function is slightly asymmetric towards lower velocities, suggesting that the known companion may be responsible for the offset from the predicted velocity. The third star is RHy 158. Our two velocity determinations differ by

**Table 5.** Heliocentric velocities (contd.)


| RHy   | MJD        | $V_{hel}$<br>kms $^{-1}$ | $\epsilon_V$<br>kms $^{-1}$ | $V^P$<br>kms $^{-1}$ | $\delta V_{hel}^P$<br>kms $^{-1}$ | $\delta V/\epsilon_V$ | $\Delta V_{epoch}$<br>kms $^{-1}$ | H $\alpha$ EW<br>Å |
|-------|------------|--------------------------|-----------------------------|----------------------|-----------------------------------|-----------------------|-----------------------------------|--------------------|
| 403   | 50000+     |                          |                             |                      |                                   |                       |                                   |                    |
|       | 793.45354  | 42.96                    | 0.5                         | 40.11                | 1.75                              | 3.5                   |                                   | 5.2                |
|       | 794.54738  | 73.76                    | 0.8                         |                      | 33.7                              | 42                    | 30.80                             | 5.7                |
|       | 1541.19393 | 69.42                    | 0.6                         |                      | 29.3                              | 49                    | 26.46                             | 7.8                |
|       | 1541.24538 | 75.41                    | 0.5                         |                      | 35.3                              | 71                    | 32.45                             | 7.5                |
|       | 1541.29500 | 78.83                    | 0.5                         |                      | 38.7                              | 77                    | 35.87                             | 6.2                |
|       | 1541.35213 | 79.99                    | 0.8                         |                      | 39.9                              | 50                    | 37.03                             | 8.1                |
|       | 1541.40799 | 78.46                    | 0.5                         |                      | 38.4                              | 77                    | 35.50                             | 6.4                |
|       | 1541.45265 | 75.15                    | 0.6                         |                      | 35.0                              | 58                    | 32.19                             | 5.9                |
|       | 1541.48272 | 72.16                    | 0.5                         |                      | 32.1                              | 64                    | 29.20                             | 3.9                |
|       | 1542.19549 | 20.23                    | 0.5                         |                      | -19.8                             | -40                   | -22.69                            | 7.0                |
|       | 1542.24438 | 27.69                    | 0.5                         |                      | -12.4                             | -25                   | -15.27                            | 5.9                |
|       | 1542.29250 | 37.20                    | 0.5                         |                      | -2.91                             | -6                    | -5.76                             | 7.2                |
|       | 1542.34197 | 47.04                    | 0.6                         |                      | 6.9                               | 12                    | 4.08                              | 6.4                |
|       | 1542.39357 | 56.73                    | 0.5                         |                      | 16.6                              | 33                    | 13.77                             | 6.2                |
|       | 1542.43815 | 64.10                    | 0.5                         |                      | 24.0                              | 48                    | 21.14                             | 6.2                |
| G8-17 | 793.39242  | 36.85                    | 0.5                         | 35.96                | 0.91                              | 1.8                   |                                   | 9.8                |


**Figure 4.** H $\alpha$  emission and typical cross-correlation results for RHy 42 observations. The left panels plot data from January, 1997; the middle panel, the December, 1997 observation; the right panel, data from December, 1999. The 1997 data clearly identify RHy 42 as a double-lined binary with components of nearly equal luminosity.


only 1.8 kms $^{-1}$  ( $\sim 4\epsilon_V$ ), but this star was also observed by Terndrup et al (2000). Their datum differs significantly from both our measurements.

RHy 49, 162, 165 and 244 all have cross-correlation functions with maxima suggestive of double-lined systems, with velocity separations of 5 to 10 kms $^{-1}$  (figure 5). The first three stars are all rapid rotators, and it is possible that rapid rotation, rather than binarity, may be responsible for the anomalous structure (although RHy 49 is resolved as binary by HST). RHy 244, however, has low  $v \sin(i)$ , and the

narrow cross-correlation peak has blueward asymmetry, suggesting a contribution from a lower-luminosity companion (see, for comparison, figures 3 and 7 in Stauffer et al, 1997). We know from HST imaging that RHy 244 has a companion at  $\Delta \sim 0.1$  arcseconds,  $\delta I \sim 1.1$  magnitudes. The observed asymmetry is not inconsistent with the expected velocity difference of these known components.



**Figure 3.** Velocity measurements from December, 1999 for RHy 42 (crosses), RHy 281 (triangles) and RHy 403 (open circles). The curve matched to the last set of observations is a sine-wave with  $P=1.276$  days, amplitude  $\pm 40 \text{ km s}^{-1}$  and  $T_0=51541.3521$  days (MJD).



**Figure 5.** The central regions of the cross-correlation functions for RHy 49, 162, 165 and 244. The RHy 244 cross-correlation peak has an asymmetric blue wing, while the remaining three functions exhibit anomalous profiles.

## 5.2 The companion of RHy 403

Our observations of RHy 403 provide sufficient information to set firm constraints on the mass of the companion. If we assume a circular orbit, as suggested by the sine curve matched against the data in figure 3, then the observed maximum velocity ( $80 \text{ km s}^{-1}$ ) and  $V^P \sim 40 \text{ km s}^{-1}$  imply an orbital velocity,  $v \sin(i) = 40 \text{ km s}^{-1}$ , and a projected semi-major axis,  $a \sin(i) \sim 701,000 \text{ km}$ , or  $0.0047 \text{ A.U.}$ . The systemic mass function is given by

$$f(m) = \frac{M_2^3 \sin^3(i)}{(M_1 + M_2)^2} = \frac{(a \sin(i))^3}{P^2}$$

where  $P$  is the period, 1.275 days. Substituting for  $P$ , we derive  $f(m) = 0.0085$ .

The fact that RHy 403 is a single-lined system allows us to estimate both a lower limit and an upper limit to  $M_2$ , the mass of the unseen companion. First, we can assume that the primary contributes most of the flux, even at  $8000\text{\AA}$ . Hence, the observed absolute magnitude provides an estimate of the mass of the primary, RHy 403A. Based on  $(BC_I, (V-I))$  data from Leggett et al (1996) and the photometry given in Table 1, we derive  $M_{bol} \sim 11$ . Matched against the 600 Myr-old models computed by Burrows et al (1993), this implies a mass of  $\sim 0.15 M_\odot$ . Substituting for  $M_1$  in the mass function gives


$$M_2^3 \sin^3(i) = 1.9 \times 10^{-4} + 0.0085 M_2^2$$

Setting  $i = 90^\circ$  gives a lower limit to the mass of the companion,  $M_2 > 0.062 M_\odot$ , the first solution of the cubic equation.

In principle, RHy 403 could have an evolved companion; in practice, we can effectively rule out this option. With an age of 625 Myrs, the lowest luminosity white dwarfs in the Hyades have  $M_V \sim 12$  and  $T_{eff} \sim 15,000\text{K}$ . Unresolved Hyades M dwarf/white dwarf binaries therefore have unusual colours, as demonstrated by HZ 9 ( $M_2 + \text{DA}$ :  $M_V = 10.6$ ,  $(V-I)=1.8$ ). A neutron star companion,  $M_2 = 1.4$  to  $2 M_\odot$ , requires  $10^{9.5} > i > 9.3$  and  $0.026 < a < 0.029 \text{ AU}$ ; the extensive mass loss during a supernova excludes the possibility of a primordial binary at these separations, while a system forming post-SN would still be expected to be bright at X-ray wavelengths. Stern et al (1994) do not detect a ROSAT source close to RHy 403 in their survey of the cluster.

Given that the companion is either a low-luminosity main-sequence star or a brown dwarf, we can use the non-detection in both the direct spectrum and, more stringently, the cross-correlation analysis, to set an upper limit to  $M_2$ . A  $0.06 M_\odot$  brown dwarf is predicted to have a temperature of  $\sim 1900\text{K}$  at age 625 Myrs, corresponding to a spectral type of L2 and a luminosity of  $10^{-4} L_\odot$ . With an absolute magnitude of  $M_I \sim 15$ , over 4 magnitudes fainter than RHy 403A, such an object lies well below our detection limit.

How faint a companion could we expect to detect? We can address this question in two ways. First, we have modelled an unequal-mass, late-type M-dwarf binary by combining appropriate spectra spanning a range of flux ratio. Based on those tests, we would expect to detect anomalies in the cross-correlation function for flux ratios exceeding 4:1 at  $\lambda > 8000\text{\AA}$ . Second, we can use observations of known spectroscopic binaries to infer detection limits. The data



**Figure 6.** The cross-correlation peak for Gl 831AB matched against our template, Gl 447. The peak is asymmetric towards higher velocities. We interpret the asymmetry as due to the known companion, Gl 831B.

plotted in figure 5 suggest that we have detected RHy244B,  $\delta I = 1.1$  magnitude, a flux ratio of  $\sim 3 : 1$ . Similarly, Gl 831 is a known binary, with  $M_V(A) = 12.5$ ,  $M_V(B) = 15.5$ ; masses,  $M_A = 0.25M_\odot$ ,  $M_B = 0.1M_\odot$ ;  $a \sim 0.21$  arcseconds; and  $P \sim 1.92$  years (Henry, 1990). Cross-correlating our spectra of Gl 831 against data for Gl 447 reveals an asymmetric peak (figure 6). Given  $\Delta M_V \sim 3$  magnitudes and  $\Delta M_K \sim 1.3$  magnitudes for the two components, the likely magnitude difference at I is close to 2 magnitudes, or a flux ratio of  $\sim 6:1$ . Our data for RHy 403 have lower signal to noise, but it is unlikely that a companion with a flux ratio exceeding 4:1 would have escaped detection.

Our flux limit corresponds to an absolute magnitude of  $M_I = 12.3$ . Matched against the Burrows et al (1993) models, this implies an upper mass limit of  $M_2 < 0.095M_\odot$  for RHy 403B.

The lowest-mass member of the Hyades currently known is LH0418+13 (Reid & Hawley, 1999), with  $M_I \sim 14$  and a mass of approximately  $0.08 M_\odot$ . There is a probability of 72% that the orbit of RHy 403B has  $i > 50^\circ$  and  $M < 0.08M_\odot$ ; the probability is 66% that  $i > 55^\circ$  and  $M < 0.075M_\odot$ . Thus, RHy 403B is the first strong candidate for a brown dwarf member of the Hyades cluster. Unfortunately, since  $asin(i) = 0.0047\text{AU}$  and  $r = 53$  parsecs, the maximum expected angular separation (for  $\frac{M_2}{M_1} \sim 2.5$ ) is only 0.25 mas. High-resolution spectroscopy at near-infrared wavelengths may provide stronger constraints on the upper mass limit.

### 5.3 The binary fraction of late-type Hyades M dwarfs

Prior to our observations, ten of the fifty-one Hyades stars listed in Table 1 were known to be in binary systems. Our spectroscopic data add three definite new systems, RHy 42, 158 and 403, with RHy 281 also a strong candidate binary. In addition, RHy 162 and 165 show indications of duplicity, although the high intrinsic rotation of these stars render less certain the interpretation of the cross-correlation data. RHy 244, 377 and, perhaps, 49 are also possible spectroscopic binaries, but all three were identified previously as binaries based on the HST imaging. Our present radial velocity data are insufficient to determine whether the spatially-resolved and hypothetical spectroscopic components are one and the same.

Forty-eight of the fifty-one M dwarfs listed in Table 1 have absolute magnitudes  $M_V > 12$ . Amongst the 47 systems (RHy 240A and B are observed separately) at least 11, and perhaps as many as 14, are binary or multiple systems - a multiplicity fraction of 23 to 30%, with a formal uncertainty of 7%. In comparison, the binary fraction amongst the (more completely surveyed) M dwarfs in the 8-parsec sample is 33% (Reid & Gizis, 1997a).

Further analysis is limited by the small numbers in both the Hyades and field M dwarf samples. However, Table 6 compares the distribution of projected separations,  $\Delta$ , adopting  $\Delta = 0.8a$  for nearby systems with known orbits. We list statistics for all 8-parsec binaries with M-dwarf primaries and for the subset where the primary has  $M < 0.3M_\odot$ . Within the admittedly large uncertainties, the field star and cluster distributions are identical.

## 6 CHROMOSPHERIC ACTIVITY IN LOW-MASS HYADES DWARFS

### 6.1 Rotation and activity

Table 8 lists the rotational velocities derived for the Hyades dwarfs in our sample. All are chromospherically active, with substantial  $H\alpha$  emission. As indicated by the individual measurements listed in Table 5, the level of activity can vary by up to 30% on relatively short timescales. This intrinsic variability is the dominant source of uncertainty in determining stellar activity from one or two observations, as is the case here. Figure 7 illustrates this effect, comparing successive observations of RHy 403 from Dec 29, 1999. Note the central reversal in the stronger  $H\alpha$  line profile. As Stauffer et al point out, the majority of slow rotators ( $v \sin(i) < 12 \text{ km s}^{-1}$ ) have similar profiles, while stars with faster rotation have either flat-topped or peaked  $H\alpha$  lines.

We are interested in determining the extent of any correlation between the level of chromospheric activity and the underlying stellar rotation. The standard paradigm (Stauffer et al, 1997) envisions a direct correlation, with activity increasing with increasing rotation until a “saturation” velocity is achieved, where the chromosphere and coronal loops are fully active. Recent observational results, however, have raised serious questions as to whether this scenario is maintained in low-mass, late-type M dwarfs (Hawley et al, 1999).


The standard rotation/activity hypothesis rests on an extension of the observed behaviour of G and K dwarfs,

**Table 6.** Distribution of separation of binary components

|              | Hyades |     | Field | all M | Field | $M < 0.3M_{\odot}$ |
|--------------|--------|-----|-------|-------|-------|--------------------|
| $\Delta$     | N      | %   | N     | %     | N     | %                  |
| $\leq 10$ AU | 7      | 50% | 17    | 50%   | 11    | 65%                |
| 10 - 100 AU  | 5      | 36  | 11    | 32    | 6     | 35                 |
| $> 100$ AU   | 1      | 14  | 6     | 18    | 0     | 0                  |

**Table 7.** Rotation and activity in low-mass Hyades stars

| Name           | v sin(i) | $M_{bol}$ | Log ( $L_{\alpha}/L_{bol}$ ) | Name        | v sin(i) | $M_{bol}$ | Log ( $L_{\alpha}/L_{bol}$ ) |
|----------------|----------|-----------|------------------------------|-------------|----------|-----------|------------------------------|
| RHy 9          | 3.5      | 10.55     | -3.88                        | RHy 23      | 4        | 10.08     | -3.86                        |
| 42             | 3.5      | 10.19     | -3.99                        | 46          | 5        | 10.28     | -3.89                        |
| 49             | 22       | 9.81      | -3.80                        | 60          | 19       | 9.66      | -3.95                        |
| 64             | 19       | 10.68     | -4.01                        | 83          | 8        | 11.29     | -3.98                        |
| 98             | 23       | 10.68     | -3.87                        | 101         | 16       | 10.89     | -4.14                        |
| 115            | 7        | 10.05     | -3.87                        | 119         | 7.5      | 10.73     | -3.98                        |
| 126            | 21       | 10.36     | -3.82                        | 129         | 3.5      | 9.94      | -3.85                        |
| 143            | 4        | 10.15     | -3.85                        | 158         | 11       | 10.14     | -3.92                        |
| 162            | 35       | 10.59     | -3.77                        | 163         | 19       | 9.83      | -3.94                        |
| 165            | 17       | 11.67     | -3.99                        | 182         | $< 3$    | 9.80      | -3.87                        |
| 199            | 16       | 10.46     | -3.99                        | 200         | 7.5      | 10.06     | -3.84                        |
| 202            | 2.5      | 10.53     | -4.04                        | 219         | 7        | 10.55     | -3.76                        |
| 221            | 2.5      | 10.50     | -3.98                        | 230         | 16       | 11.47     | -3.91                        |
| 231            | 6.5      | 9.69      | -3.73                        | 240A        | 14.5     | 11.68     | -3.90                        |
| 240B           | 14.5     | 11.72     | -4.09                        | 242         | 6        | 10.01     | -3.93                        |
| 244            | 2.5      | 9.46      | -3.76                        | 260         | 2.5      | 10.77     | -3.90                        |
| 281            | 9.5      | 11.32     | -3.66                        | 297         | 13       | 10.52     | -3.80                        |
| 298            | 6        | 10.10     | -3.86                        | 301         | 12       | 10.57     | -3.72                        |
| 331            | 6.5      | 10.66     | -3.94                        | 346         | 3.5      | 10.44     | -3.98                        |
| 367            | 3        | 10.11     | -3.82                        | 369         | $< 2.5$  | 10.05     | -3.99                        |
| 371            | 14       | 9.23      | -3.90                        | 376         | 4        | 10.00     | -3.83                        |
| 377            | 9        | 9.79      | -3.90                        | 386         | 9        | 12.00     | -3.95                        |
| 390            | 18       | 11.25     | -3.77                        | 391         | 24       | 9.94      | -3.85                        |
| 399            | $< 2.5$  | 9.64      | -3.89                        | 402         | 11.5     | 10.94     | -3.75                        |
| 403            | 5.5      | 11.10     | -3.82                        |             |          |           |                              |
| <hr/>          |          |           |                              |             |          |           |                              |
| Jones et al    |          |           |                              |             |          |           |                              |
| Br 262         | 37       | 11.40     | -4.08                        | Br 804      | 31       | 11.81     | -3.86                        |
| <hr/>          |          |           |                              |             |          |           |                              |
| Stauffer et al |          |           |                              |             |          |           |                              |
| RHy 26         | $< 6$    | 8.96      | -3.76                        | RHy 28      | 11.5     | 9.45      | -3.87                        |
| 62             | $< 6$    | 8.28      | -4.03                        | 99          | $< 6$    | 9.03      | -3.89                        |
| 100            | $< 6$    | 8.93      | -4.35                        | 123         | $< 6$    | 8.69      | -5.09                        |
| 142            | 11.5     | 8.42      | -3.83                        | 161         | $< 6$    | 9.35      | -3.88                        |
| 189            | $< 6$    | 9.38      | -4.66                        | 203         | $< 6$    | 9.09      | -5.05                        |
| 211            | 6        | 8.62      | -4.39                        | 269         | 26       | 8.79      | -3.75                        |
| 283            | $< 6$    | 8.53      | -3.93                        | 289         | 6        | 9.07      | -4.71                        |
| 294            | 11       | 8.79      | -3.83                        | 296         | $< 6$    | 9.33      | -3.87                        |
| 322            | $< 6$    | 9.52      | -3.35                        | 349         | $< 6$    | 9.11      | -3.83                        |
| 350            | $< 6$    | 10.16     | -3.40                        | 384         | 25       | 8.52      | -3.70                        |
| GH 7-33        | 12.5     | 8.48      | -3.74                        | Lei 44      | 18       | 8.61      | -3.63                        |
| Lei 73         | $< 6$    | 8.72      | -4.86                        | LP 357-160  | $< 6$    | 8.73      | -4.67                        |
| LP 358-716     | $< 6$    | 8.93      | -4.05                        | LP 359-2423 | 9.5      | 8.79      | -3.76                        |
| LP 359-262     | $< 6$    | 8.84      | -4.55                        | LP 413-93   | $< 6$    | 9.13      | -4.14                        |
| LP 535-101     | $< 6$    | 9.09      | -4.10                        | vA 54       | $< 6$    | 8.94      | -3.94                        |
| vA 106         | 10       | 9.16      | -3.82                        |             |          |           |                              |
| <hr/>          |          |           |                              |             |          |           |                              |
| Terndrup et al |          |           |                              |             |          |           |                              |
| RHy 128        | 19       | 10.88     | -3.86                        | RHy 138     | 22       | 11.66     | -3.96                        |
| 206            | 24       | 10.47     | -3.93                        | LP 416-573  | 6        | 10.18     | -4.01                        |
| LP 358-735     | 25.5     | 10.16     | -3.82                        | LP 476-648  | 6        | 10.18     | -3.94                        |
| LP 358-717     | 18       | 10.64     | -3.86                        |             |          |           |                              |



**Figure 8.** Chromospheric activity and rotation in Hyades dwarfs. The upper panel (a) plots activity,  $\log L_\alpha / L_{bol}$ , for M dwarfs from this paper (circles), and from previous studies (Stauffer et al (1997), Jones et al (1996), Terndrup et al (2000) - all triangles) Solid points mark known binaries. The middle panel (b) plots the ( $M_{bol}$ ,  $v \sin(i)$ ) relation for the same stars, using the same symbols. The lowest panel (c) plots rotation against activity, where all of the Hyades observations are divided into three subsets based on  $M_{bol}$ . Solid points mark stars with  $M_{bol} < 10$ ; open circles have  $10 < M_{bol} < 11$ ; the faintest stars are marked as crosses.

where there is a clear activity-rotation correlation. Both parameters are believed to be age dependent, with the star spinning down as it loses angular momentum due to coupling of the star's magnetic field with the local medium (Skumanich, 1972). Since activity in these stars is powered by a shell dynamo (Parker, 1955), located at the boundary between the radiative core and the outer convective layers, the activity-rotation correlation is expected. This mechanism, however, cannot be present in fully convective low-mass stars. Indeed, early studies predicted a break in the level of activity at a mass of  $\sim 0.3M_\odot$ , where the radiative core disappears. The continued presence of high levels of activity in these stars is probably due to the presence of a turbulent (shear) dynamo mechanism (Durney et al, 1993). In that case, there is no expectation of a direct correlation with rotational velocity.

Investigating this question requires a reliable measurement of the level of activity in stars with temperatures between 4000 and 2400K. The equivalent width of the H $\alpha$  emission line is frequently used for this purpose. This parameter, however, provides a misleading measure of activ-


ity, since the equivalent width produced by a given line flux depends on the level of the underlying continuum. This is a particularly important consideration for M dwarfs, since the R<sub>C</sub>-band flux decreases rapidly with decreasing temperature from spectral type M0 ( $\sim 4000$ K) through M5 ( $\sim 2800$ K) to M9 ( $\sim 2100$ K). This complication can be avoided by calculating the parameter  $L_\alpha / L_{bol}$ , the fraction of the total luminosity emitted in H $\alpha$ . This provides a measure of stellar activity *independent* of both temperature and luminosity.

We have computed bolometric magnitudes for the stars in our sample using the (V-I) colours and the relation

$$BC_I = 0.02 + 0.575(V - I) - 0.155(V - I)^2$$

derived from Leggett et al's (1996) data for field M dwarfs. Table 7 lists those magnitudes, together with data for Hyades stars from Stauffer et al (1997), Jones et al (1996) and Terndrup et al (2000). Deriving  $L_\alpha$  for these stars requires an estimate of the continuum flux at 6563Å,  $F_C$ . Reid et al (1995b) demonstrated that  $F_C$  can be derived from the R<sub>C</sub>-band flux,  $F_R$ , from the relation

$$F_C = 1.45F_R$$



**Figure 9.** H $\alpha$  emission and typical cross-correlation results for Activity/rotation correlations for Pleiades stars. The data are taken from Terndrup et al (2000), Basri & Marcy (1995) and Jones et al (1996). Pleiades stars are plotted as solid points; all Hyades stars from figure 8 are plotted as crosses for comparison.

All of the stars in Table 1 have R-band photometry (Reid, 1993 and references therein). For stars with multiple observations, the H $\alpha$  fluxes are derived using the average equivalent width (with the exception of the MJD 794.46 observation of RHy 281, discussed further below). Variability, as noted above, leads to typical uncertainties of  $\pm 15\%$ .

The uppermost panel in figure 8 shows the ( $M_{bol}$ ,  $\log(L_{\alpha}/L_{bol})$ ) distribution for Hyades M dwarfs. Contrary to the suggestion by Terndrup et al (2000), There is no evidence for a decline in activity with decreasing luminosity. Known binaries and single stars are distinguished using different symbols on the diagram, but there is no evidence that the former are more (or less) active than the latter. These data confirm results derived previously by Reid et al (1995b), a paper which unaccountably seems to have escaped Terndrup et al’s attention.


Figure 8b plots the ( $v \sin(i)$ ,  $M_{bol}$ ) distribution for the Hyades stars, illustrating the well known tendency towards higher rotation with decreasing mass. However, figure 8c shows that there is no correlation between the level of chromospheric activity and rotation: a Hyades M dwarf with  $L_{\alpha}/L_{bol} \sim 1.5 \times 10^{-4}$  can have a measured rotational ve-

locity anywhere between  $< 2$  and  $40 \text{ km s}^{-1}$ , the full range measured for cluster stars.

This result is in good agreement with data for late-type field M dwarfs (Hawley et al, 1999), but directly contradicts Terndrup et al’s analysis of a subset of the present sample. The origin of the discrepancy rests with Terndrup et al’s use of equivalent width as an estimator of stellar activity. As noted above (and, indeed, by Terndrup et al), the equivalent width of an emission line depends on both the total line flux and the level of the local continuum. Thus, early-type Hyades M dwarfs have weaker emission than late-type dwarfs, even though the chromosphere is at the same *proportional* level of activity. Further, figure 8b shows that late-type dwarfs (with higher H $\alpha$  EW) tend to have more rapid rotation. Thus, the correlation between “activity” and rotation claimed by Terndrup et al is actually the (mass/ $M_{bol}$ ,  $v \sin(i)$ ) correlation illustrated in figure 8b.

Terndrup et al also present rotation and activity measurements for low-mass Pleiades stars. Figure 9 plots their data, together with observations from Jones et al (1996) and





**Figure 10.** H $\alpha$ , He I 6678Å and Ca II 8500Å emission during the flare outburst of RHy 281

Basri & Marcy (1995), using  $L_{\alpha}/L_{bol}$  rather than H $\alpha$  EW<sup>†</sup>. With an age of  $\sim 120$  Myrs, the Pleiades stars are more luminous (and hotter) at a given mass, and significantly more active. Thus, the mass range is  $\sim 0.1 < \frac{M}{M_{\odot}} < 0.6$ , similar to the Hyades stars. While the measured rotational velocities approach  $100 \text{ km s}^{-1}$ , there is no evidence for any correlation between rotation and the level of chromospheric activity.

These results are consistent with Hawley et al's (1999) proposal that the mechanism driving chromospheric activity changes as one moves from early-type M dwarfs to mid- and late-type dwarfs. As convection becomes increasingly important and the radiative core shrinks, the shell dynamo weakens. In contrast, the turbulent dynamo increases in strength, supplanting the shell dynamo entirely in fully convective stars. As a result, rotation is irrelevant to the level of activity present in dwarfs of spectral type M3 and later.

## 6.2 The flare on RHy 281

One observation in Table 5 stands out: on December 12, 1997, the measured equivalent width of H $\alpha$  emission in RHy


281 is  $\sim 23\text{\AA}$ . As figure 10 shows, we happened to catch the star during a flare outburst. In addition to H $\alpha$  emission, He I 6678Å is present with an equivalent width of  $1.15\text{\AA}$  core emission is detectable in the K I 7665/7699 doublet, and we measure equivalent widths of  $0.85$  and  $0.35 \text{\AA}$  for the 8500 and 8663 Å components of the Ca II near-infrared triplet.

Since our observations concentrated almost exclusively on stars within a narrow range of absolute magnitude,  $11.5 < M_V < 16$ , we can derive an approximate estimate of the frequency of flare outbursts in these stars. In total, we expended  $\sim 67,000$  seconds integrating on M dwarfs in the course of this project; the RHy 281 flare was detected in an 1800 second exposure, and we assume that the flare persisted at a detectable level for the duration of the exposure. Given that assumption, we deduce that low-mass Hyades dwarfs are in outburst for  $\sim 2.5\%$  of the time. This compares with an upper limit of  $< 7\%$  for the duty cycle estimated for the M9 field dwarf, BRI0021 (Reid et al, 1999).

## 7 SUMMARY AND CONCLUSIONS

We have obtained high resolution spectroscopy of 51 low-mass dwarfs in the Hyades cluster. At least three are confirmed as spectroscopic binaries: two double-lined systems,

<sup>†</sup> Note that using (V-I) instead of  $M_{bol}$  in these diagram does not change the overall morphology



**Figure 7.**  $H\alpha$  line profiles from successive observations of RHy 403 (MJD 1541.245 and 1541.295).

RHy 42 and 244, the latter previously identified as a binary in HST imaging; and RHy 403, a short-period single-lined system with an extremely low-mass secondary. Based on the mass function for the last star and the absence of any detection of the secondary at  $8000\text{\AA}$ , we infer mass limits of  $0.07 \leq \frac{M}{M_{\odot}} \leq 0.095$ , making this unseen companion a strong brown dwarf candidate.

Four other stars are possible binaries: RHy 158 shows strong indications of velocity variability; RHy 281 lies  $\sim 6 \text{ km s}^{-1}$  from the expected velocity of Hyades members; RHy 221 and 377 are also offset in velocity, although to smaller extent. Three further stars have unusual cross-correlation functions, although it is not clear whether these stem from binarity or rapid rotation. Combining the new discoveries with previous HST and ground-based identifications, we derive a binary fraction for Hyades M dwarfs of 23 to 30 %, similar to results derived from nearby field M dwarfs.

We have also used our data to determine rotational velocities and explore possible correlations between chromospheric activity and rotation. Combined with literature data, we find that the level of chromospheric activity, as measured by the fractional luminosity emitted in the  $H\alpha$  line,  $L_{\alpha}/L_{\text{bol}}$ , is nearly constant for M dwarfs with  $8 < M_{\text{bol}} < 12$ . There is no evidence for a significant correlation between activity and rotation as measured by  $v \sin(i)$ . We interpret this as confirmation that the rotational dynamo is not present in fully convective mid- and late-type M dwarfs. Activity is probably driven by a shear dynamo mechanism, as proposed by Hawley et al (1999).

## Acknowledgements

Funding for SM was provided by STScI Grant GO-08146.01-97A. INR would like to thank Wal Sargent for the allocation of Keck time in December, 1999. This analysis is based on observations obtained at the W. M. Keck Observatory, which is operated by the California Association for Research in Astronomy, a partnership of the University of California, the California Institute of Technology and the National Aeronautics and Space Administration. The observation was made possible by a generous grant from the W. M. Keck Foundation.

## REFERENCES

- Basri, G., Marcy, G.W. 1995, AJ, 109, 762  
 Burrows, A., Hubbard, W.B., Saumon, D., Lunine, J.I. 1993, ApJ, 406, 158  
 Cochran, W.D., Hatzes, A.P. 1999, DPS, 31, 0902  
 Delfosse, X., Forveille, T., Perrier, C., Mayor, M. 1998, A & A, 331, 581  
 Duquennoy, A. & Mayor, M. 1991, A&A, 248, 485  
 Duney, B.R., De Young, D.S., Roxburgh, I. 1993 Sloar Physics, 145, 207  
 Fischer, D.A. & Marcy, G.W. 1992, ApJ, 396, 178  
 Giclas, H.L., Burnham, R., Thomas, N.C., 1972, The Lowell Observatory Proper Motion Survey, Lowell Obs. Bull. 158  
 Gizis, J.E., Reid, I.N. 1995, AJ, 110, 1248  
 Gray, D.F., 1992, in The Observation and Analysis of Stellar Photospheres (New York: Cambridge univ. Press)  
 Griffin, R.F., Gunn, J.E., Zimmerman, B.A. & Griffin, R.E.M. 1988, AJ, 96, 172  
 Hagen, H.-J., Groote, D., Engels, D., Reimers, D. 1995, A & AS, 111, 195  
 Hanson, R.B. 1975, AJ, 80, 379  
 Hawley, S.L., Reid, I.N., Gizis, J.E. 1999, in Workshop on Cool Stars and Giant Planets, ed. C. Griffiths & M. Marley, ASP Conf. Ser. No. 214,  
 Henry, T.J. 1990, Ph.d. thesis, University of Arizona  
 Henry, T.J., McCarthy, D.W. 1993, AJ, 106, 773  
 Jones, B.F., Fischer, D.A., Stauffer, J.R. 1996, AJ, 112, 1562  
 Leggett, D.K., 1992, ApJS, 82, 351  
 Leggett, S.K., Harris, H.C. & Dahn, C.C. 1994, AJ, 108, 944  
 Leggett, S.K., Allard, F., Berriman, G., Dahn, C.C., Hauschildt, P. 1996, ApJS, 104, 117  
 Marcy, G.W., Benitz, K.J. 1989, ApJ, 344, 441  
 Marcy, G.W., Butler, R.P. 1992, PASP, 104, 270  
 Parker, E.N., 1955, ApJ, 122, 293  
 Patience, J., Ghez, A.M., Reid, I.N., Weinberger, A.J., Matthews, K. 1998, AJ, 115, 1972  
 Perryman MAC, Brown AGA, Lebreton Y, Gómez A, Turon C, Cayrel de Strobel G, Mermilliod J-C, Robichon N, Kovalevsky J, Crifo F 1998. A&A, 331, 81  
 Reid, I.N. 1992, MNRAS, 257, 257  
 Reid, I.N. 1993, MNRAS, 265, 785  
 Reid, I.N., Hawley, S.L., Gizis, J.E. 1995a, AJ, 110, 1838  
 Reid, I.N., Hawley, S.L., Mateo, M. 1995a, MNRAS, 272, 828  
 Reid, I.N., Gizis, J.E. 1997a, AJ, 113, 2246  
 Reid, I.N., Gizis, J.E. 1997b, AJ, 114, 1992  
 Reid, I.N., Hawley, S.L. 1999, AJ, 117, 343  
 Reid, I.N., Kirkpatrick, J.D., Gizis, J.E., Liebert, J. 1999, ApJ, 527, L105  
 Sandage, A., Kowal, C. 1986, AJ, 91, 1140  
 Skumanich, A., 1972, ApJ, 171, 565  
 Stauffer, J.R., Balachandran, A.C., Krishnamurthi, A., Pinsonneault, M., Terndrup, D.M., Stern, R.A. 1997, ApJ, 475, 604

- Stern, R. A., Schmitt, J.H.M.M, Pye, J.P., Hodgkin, S.T., Stauffer, J.R., Simon, T. 1994, ApJ, 427, 808
- Terndrup, D.M., Stauffer, J.R., Pinsonneault, M.H., Sills, A., Yuan, Y., Jones, B.F., Fiscer, D., Krishnamurthi, A. 2000, AJ, 119, 1303
- Tonry, J., Davis, M. 1979, AJ, 84, 1511
- Vogt, S.S. et al., 1994, S.P.I.E., 2198, 362.